

学校编码: 10384

密级_____

学号: 22620060153352

厦 门 大 学

博 士 学 位 论 文

中国典型海区海洋古菌生态功能及环境适应机制的研究

Ecological Function and Environmental Adaptation
Mechanisms of Pelagic Archaea in Typical Areas of China
Seas

胡安谊

指导教师姓名: 焦念志 教授

专 业 名 称: 环 境 科 学

论文提交日期: 2010 年 6 月

论文答辩日期: 2010 年 9 月

学位授予日期: 2010 年 月

2010 年 9 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

（ ） 2. 不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目 录

摘 要.....	I
ABSTRACT.....	III
第一章 前言	1
1.1 海洋微生物生态学研究的发展历程.....	1
1.2 海洋古菌研究的发展历程及研究现状.....	3
1.2.1 古菌简介.....	3
1.2.2 海洋古菌	6
1.2.3 氨氧化古菌	14
1.2.4 浮游泉古菌自养固碳	25
1.3 本论文的设计与研究意义.....	28
第二章 中国典型海区浮游古菌分子生态学研究.....	31
2.1 研究背景	32
2.2 材料和方法	33
2.2.1 采样和环境参数的测定.....	33
2.2.2 基因组 DNA 提取, PCR 扩增, 克隆文库构建	34
2.2.3 扩增产物 RFLP 及多样性分析	36
2.2.4 系统发育分析.....	36
2.3 结果.....	37
2.3.1 采样站位的环境参数.....	37
2.3.2 古菌 16S rRNA 基因克隆文库 RFLP 分析	38
2.3.3 古菌 16S rRNA 基因序列系统发育分析.....	40
2.4 讨论.....	41
2.4.1 中国海区古菌群落结构.....	41
2.4.2 系统发育分析.....	44
第三章 南海 MD2896 沉积物柱状样古菌群落结构分析	45
3.1 研究背景	46
3.2 材料与方法	47
3.2.1 采样和环境参数的测定.....	47
3.2.2 基因组 DNA 提取, PCR 扩增和克隆文库构建	48

3.2.3 扩增产物 RFLP 及多样性分析	49
3.2.4 系统发育分析	50
3.2.5 统计分析	50
3.3 结果	50
3.3.1 南海 MD2896 站沉积物柱状样化学参数	50
3.3.2 古菌 16S rRNA 基因多样性	51
3.3.3 古菌 16S rRNA 基因系统发育分析	51
3.3.4 古菌 16S rRNA 克隆文库群落统计分析	54
3.3.5 MD2896 站表层沉积物中的古菌 <i>amoA</i> 基因	55
3.4 讨论	56
3.4.1 南海 MD2896 站沉积物柱状样中古菌群落结构	56
3.4.2 南海 MD2896 站表层沉积物中古菌的生态功能	58
第四章 南海古菌群落结构与生态功能在深度尺度上的变迁	60
4.1 研究背景	61
4.2 材料和方法	62
4.2.1 采样站位及环境参数测定	62
4.2.2 基因组 DNA 提取	64
4.2.3 定量 PCR(qPCR)分析	64
4.2.4 扩增产物 DGGE 分析	67
4.2.5 泉古菌 <i>amoA</i> 和 <i>accA</i> 基因文库构建以及测序分析	68
4.2.6 统计分析	68
4.3 结果	69
4.3.1 采样站位的水团性质	69
4.3.2 古菌 16S rRNA、 <i>amoA</i> 和 <i>accA</i> 基因以及 AOB 的 <i>amoA</i> 基因丰度	69
4.3.3 不同基因之间的线性回归分析	73
4.3.4 颗粒粘附古菌的 16S rRNA 基因丰度	73
4.3.5 古菌 16S rRNA 基因 DGGE 图谱分析	74
4.3.6 泉古菌 <i>amoA</i> 和 <i>accA</i> 基因克隆文库分析	76
4.4 讨论	78
4.4.1 南海古菌各个类群的丰度	78
4.4.2 深度梯度上泉古菌生态功能的变迁	81
4.4.3 海洋泉古菌的适应和进化演变	83

4.5 小结	85
第五章 东海真光层和弱光层上层浮游泉古菌的生态位分化现象及机制	86
5.1 研究背景	87
5.2 材料和方法	88
5.2.1 采样站位及环境参数测定	88
5.2.2 基因组 DNA 提取	88
5.2.3 定量 PCR(qPCR)分析	88
5.2.4 细菌 16S rRNA 基因 T-RFLP 及泉古菌 16S rRNA DGGE 分析	89
5.2.5 泉古菌 <i>amoA</i> 和 <i>accA</i> 基因文库构建以及测序分析	90
5.2.6 统计分析	90
5.3 结果	90
5.3.1 采样站位地理位置及理化参数	90
5.3.2 古菌 16S rRNA, <i>amoA</i> 和 <i>accA</i> 基因丰度	91
5.3.3 细菌、泉古菌 16S rRNA 基因遗传多样性分析	93
5.3.4 泉古菌 <i>amoA</i> 和 <i>accA</i> 基因克隆文库的多样性及系统发育分析	95
5.3.5 泉古菌 <i>amoA</i> 和 <i>accA</i> 基因生态型模拟及群落图谱分析	96
5.4 讨论	98
5.4.1 海洋上层水体中泉古菌的生态功能	98
5.4.2 海洋泉古菌的进化与生态适应机制	100
第六章 青藏高原湖泊氨氧化古菌和氨氧化细菌的群落结构及与环境因子之间的联系	102
6.1 研究背景	103
6.2 材料和方法	104
6.2.1 采样站位及环境参数测定	104
6.2.2 流式细胞术及叶绿素(Chl <i>a</i>)分析	105
6.2.3 基因组 DNA 提取, PCR 扩增, <i>amoA</i> 基因文库构建以及测序分析	106
6.2.4 多样性指数计算以及统计分析	107
6.3 结果	108
6.3.1 湖泊环境参数	108
6.3.2 青藏高原湖泊古菌和细菌 <i>amoA</i> 基因的多样性分析	108
6.3.3 古菌 <i>amoA</i> 基因的系统发育分析	111

6.3.4 细菌 <i>amoA</i> 基因的系统发育分析	112
6.3.5 氨氧化微生物群落结构聚类分析.....	112
6.3.6 环境因子与 AOA 群落结构之间的联系.....	114
6.4 讨论.....	115
6.4.1 青藏高原湖泊中 AOA 和 AOB 的多样性和分布	115
6.4.2 青藏高原湖泊 AOA 和 AOB 的系统发育分析	116
6.4.3 氨氧化微生物群落结构对环境因子的响应.....	118
6.5 小结.....	119
第七章 本论文的主要结论和创新点.....	121
7.1 主要结论.....	121
7.2 创新点.....	122
参 考 文 献	123
攻读博士学位期间完成的论文	156
致 谢.....	157

Contents

ABSTRACT (In Chinese).....	I
ABSTRACT (In English)	III
Chapter 1 Preface.....	1
1.1 Major advances in microbial oceanography	1
1.2 Major advances and current research status of marine archaea.....	3
1.2.1 Brief introduction about archaea.....	3
1.2.2 Marine archaea.....	6
1.2.3 Ammonia-oxidizing archaea.....	14
1.2.4 Autotrophic carbon fixation in planktonic <i>Crenarchaeota</i>	25
1.3 Design of this thesis and research significance	28
Chapter 2 Molecular ecological study on planktonic archaea in China Seas	31
2.1 Background.....	33
2.2 Materials and Methods	33
2.2.1 Sampling and determination of environmental variables	33
2.2.2 DNA extraction, PCR amplification and clone library construction	34
2.2.3 PCR-RFLP screening of clone libraries and diversity statistics	36
2.2.4 Phylogenetic analysis.....	36
2.3 Results.....	37
2.3.1 Environmental variables in the sampling sites.....	37
2.3.2 RFLP analysis of archaeal 16S rRNA gene libraries.....	38
2.3.3 Phylogenetic analysis of archaeal 16S rRNA genes	40
2.4 Discussion	41
2.4.1 Comparison of archaeal community structures in China Seas.....	41
2.4.2 Implication of observed phylogenetic groups.....	44
Chapter 3 Community structure of archaea from deep-sea sediment of the South China Sea	45
3.3 Background.....	46
3.2 Materials and Methods	47
3.2.1 Sampling and determination of environmental variables	47
3.2.2 DNA extraction, PCR amplification and clone library construction	48
3.2.3 PCR-RFLP screening of clone libraries and diversity statistics	49

3.2.4 Phylogenetic analysis.....	50
3.2.5 Statistical analysis.....	50
3.3 Results.....	50
3.3.1 Analysis of environmental variables.....	50
3.3.2 Diversity of archaeal 16S rRNA genes.....	51
3.3.2 Phylogenetic analysis of archaeal 16S rRNA genes.....	51
3.3.3 Community classification of archaeal 16S rRNA gene libraries.....	54
3.3.4 Archaeal <i>amoA</i> genes in the surface sediment.....	55
3.4 Discussion	56
3.4.1 Community structure of archaea in the core sediment from station MD2896.....	56
3.4.2 Ecological function of archaea in the surface sediment.....	58
Chapter 4 Community structure and function of planktonic <i>Crenarchaeota</i>: Changes with depth in the South China Sea.....	60
4.1 Background.....	61
4.2 Materials and Methods	62
4.2.1 Sampling and determination of environmental variables	62
4.2.2 DNA extraction.....	64
4.2.3 Quantitative PCR (qPCR) analysis	64
4.2.4 DGGE analysis.....	67
4.2.5 Construction of crenarchaeal functional genes clone libraries and sequencing.....	68
4.2.6 Statistical analysis.....	68
4.3 Results.....	69
4.3.1 Properties of different water bodies.....	69
4.3.2 Abundances of 16S rRNA-, <i>amoA</i> - and <i>accA</i> genes of free-living archaea.....	69
4.3.3 Correlations among different gene abundances (<i>amoA</i> , <i>accA</i> , 16S rRNA) of free-living archaea	73
4.3.4 Abundance of 16S rRNA genes of particle-attached archaea.....	73
4.3.5 DGGE analysis of archaeal communities	74
4.3.6 Clone libraries of <i>amoA</i> - and <i>accA</i> genes of free-living archaea.....	76
4.4 Discussion	78
4.4.1 Abundance of different genes of archaea.....	78
4.4.2 Changes in ecological function of <i>Crenarchaeota</i> at different depths	81
4.4.3 Adaptability of marine <i>Crenarchaeota</i>	83

4.5 Conclusion.....	85
Chapter 5 Niche-partitioning of pelagic <i>Crenarchaeota</i> in the euphotic and upper twilight zones of the East China Sea.....	86
5.1 Background.....	87
5.2 Materials and Methods	88
5.2.1 Sampling and determination of environmental variables	88
5.2.2 DNA extraction.....	88
5.2.3 Quantitative PCR (qPCR) analysis	88
5.2.4 T-RFLP analysis of bacterial 16S rRNA genes and DGGE analysis of crenarchaeal 16S rRNA genes.....	89
5.2.5 Construction of crenarchaeal functional genes clone libraries and sequencing.....	90
5.2.6 Statistical analysis.....	90
5.3 Results.....	90
5.3.1 Site description and analysis of environmental variables	90
5.3.2 Abundances of 16S rRNA-, <i>amoA</i> - and <i>accA</i> genes of archaea	91
5.3.3 Genetic fingerprinting analyses of 16S rRNA genes of bacteria and archaea	93
5.3.4 Diversity and phylogeny of crenarchaeal <i>amoA</i> and <i>accA</i> genes	95
5.3.5 Ecotype simulation and community classification of crenarchaeal <i>amoA</i> and <i>accA</i> genes	96
5.4 Discussion	98
5.4.1 Ecological roles of pelagic <i>Crenarchaeota</i> in the upper ocean	98
5.4.2 Environmental adaptation mechanisms of marine <i>Crenarchaeota</i>	100
Chapter 6 Community structures of ammonia-oxidising archaea and bacteria in high-altitude lakes on the Tibetan Plateau .	102
6.1 Background.....	103
6.2 Materials and Methods	104
6.2.1 Sampling and determination of environmental variables	104
6.2.2 Flow cytometry and Chl <i>a</i> measurement	105
6.2.3 DNA extraction, PCR, cloning and sequencing.....	106
6.2.4 Diversity indices and statistical analyses.....	107
6.3 Results.....	108
6.3.1 Lake characteristics.....	108
6.3.2 Diversity of archaeal and bacterial <i>amoA</i> genes in Tibetan lakes	108

6.3.3 Phylogeny of archaeal <i>amoA</i> genes	111
6.3.4 Phylogeny of bacterial <i>amoA</i> genes.....	112
6.3.5 Community classification of AOA	112
6.3.6 Environmental factors explaining AOA community structures.....	114
6.4 Discussion	115
6.4.1 Diversity and distribution of AOA and AOB in Tibetan lakes.....	115
6.4.2 Phylogeny of AOA and AOB	116
6.4.3 Response of the AOA community to environmental changes	118
6.5 Conclusion.....	119
Chapter 7 Conclusions and originality of this thesis.....	121
7.1 Major conclusions.....	121
7.2 Originality	122
References	123
Manuscripts during graduate study.....	156
Acknowledgements	157

摘 要

浮游古菌是海洋微生物的重要组成部分之一。在真光层以下的深海大洋,海洋
泉古菌Marine Group I (MGI) 类群的生物量可占总原核生物量的20%。最近,环
境基因组学研究的突破性进展和海洋泉古菌MGI类群纯培养株系的获得揭示出
大部分MGI类群都是氨氧化古菌 (Ammonia-Oxidizing Archaea, AOA)。由于MGI
类群是海洋中数量最为丰富的微型生物之一,其必然在海洋碳、氮循环中起着重
要作用,目前AOA已成为环境微生物生态学的研究热点之一。虽然针对浮游古
菌尤其是MGI类群的生态调查表明该类群广泛分布于海洋水体中,但在中国海
区,仅本课题组在2002年对东海长江口区域的浮游古菌进行了初步的多样性调查
(Zeng et al., 2007)。对中国海区浮游古菌尤其是MGI类群的群落结构与生态功
能的了解还十分有限。本论文通过分子生态学手段对中国海区浮游古菌开展了以
下几方面的工作,并在对海洋泉古菌研究认知的基础上,进一步比较了湖泊泉古
菌与海洋泉古菌环境适应机制的异同,拓展了对水环境常温古菌的认识:

1. 应用古菌16S rRNA基因调查了中国东海和南海六个站位表层水体中的古菌
群落组成。研究表明河口区域的浮游古菌遗传多样性最高,其次是外海
区域,而近海水体中的浮游古菌多样性最低。系统发育分析表明,中国海区
的浮游古菌可分为泉古菌和广古菌:由于冲淡水的输入,河口的古菌群落由
浮游古菌和陆源古菌组成;近岸和外海区域则没有受到陆源输入的影响。中
国海区分布的优势浮游古菌类群在其他海区的相对丰度偏低,说明中国海区
浮游古菌的地理分布模式有着一定的地域性特色。
2. 对南海11 m长沉积物柱状样中的古菌群落研究发现,分布于该沉积物的泉古
菌和广古菌的主要系统发育类群是普遍分布于其他大洋沉积物的优势类群。
基于RFLP带型的聚类分析和基于系统发育类群的RDA分析表明该沉积物中
的古菌群落组成有深度分布趋势。古菌功能基因PCR筛选表明,表层沉积物
中的MGI类群含有氨单加氧酶 α 亚基编码基因—*amoA*基因。由于本课题组曾
在同一样品检测到厌氧氨氧化细菌的16S rRNA基因序列和氨氧化细菌的16S
rRNA基因序列,说明古菌与其他N循环相关的微型生物一起参与了南海沉积

物的N循环过程。

3. 结合古菌16S rRNA基因、泉古菌氨氧化基因*amoA*和自养固碳基因*accA*研究了南海三个远洋站位全水柱剖面的古菌群落结构与生态功能。在不同站位的相同水系中，古菌16S rRNA基因、*amoA*和*accA*基因的丰度量级和遗传多样性都较为相似；而在垂直梯度上，深层水体（> 1000 m）中的古菌基因丰度要显著低于上层（< 200 m）和中层水体（200 m - 1000 m）中的古菌基因丰度（除了古菌*accA*基因在上层水体中极为稀少）。古菌16S rRNA基因DGGE分析和功能基因克隆文库结果表明上层水体与中、深层水体的古菌种群存在明显的进化分歧。统计分析不同基因间的丰度比表明，上层水体中的MGI类群缺少自养固碳基因，深层水体的MGI类群缺少氨氧化基因。这些结果预示了浮游泉古菌在大洋不同水体有着种群演替和生态功能的变迁。
4. 结合泉古菌16S rRNA基因、*amoA*和*accA*基因研究了东海黑潮区域两个站位真光层和弱光层上层水体中的泉古菌生态位分化现象及环境适应机制。与我们在南海的研究结果相似，东海上层水体中的大部分MGI类群含有*amoA*基因，但缺少*accA*基因，说明上层大洋中的浮游泉古菌可能是以化能无机异养方式生存。生态型模拟计算表明，上层水体中的浮游泉古菌同样分化为适应不同深度梯度的生态型，说明光可能是造成这种生态位分化的重要因子。
5. 通过泉古菌*amoA*基因PCR—克隆文库方法比较了五个青藏高原湖泊的氨氧化微生物的群落结构。系统发育分析和统计分析表明湖泊之间的氨氧化古菌群落结构受盐度调控，而不同湖泊中的氨氧化细菌则隶属于*Nitrosomonas*属的一个单一的系统发育分支。与海洋浮游泉古菌在深度梯度上的生态位分化相反，湖泊内部不同水层的氨氧化古菌趋同进化，说明湖泊和海洋中的泉古菌在环境适应机制上存在着差别。

关键词：浮游古菌；泉古菌；氨氧化古菌；16S rRNA 基因；*amoA* 基因；*accA* 基因自养固碳；系统发育分析；生态功能；环境适应机制；中国海；青藏高原湖泊

Abstract

Planktonic archaea are one of the most important components of Marine microbes. Marine *Crenarchaeota* Group I (MGI), one phylogenetic group of the planktonic archaea, are typically relatively more abundant in the dark ocean and are thought to account for 20% of all prokaryotic cells in the global ocean. Recently, the great breakthrough in studies of metagenomics and pure culture of MGI revealed that functional group of them—ammonia-oxidizing archaea (AOA). Given the suggestion abundance of MGI in the marine environments, AOA would play an important role in Earth's biogeochemistry and have become a new hotspot in environmental microbial ecology. Although numerous studies have demonstrated that MGI are one of the most abundant and ubiquitous life forms in the ocean, reports about diversity and distribution of planktonic archaea in China Seas are very rare. Zeng et al. (2007) have studied the genetic diversity of planktonic archaea in the Yangtze River estuary. However, the community structure and ecological function of planktonic archaea in China Seas is still unclear. Here we investigated the geographic distribution, genetic diversity, ecological function and environmental adaptation mechanisms of planktonic archaea in typical areas of China Seas and Tibetan lakes by using molecular ecological methods.

1. The community structures of planktonic archaea were investigated for the surface water from six stations in the East (ECS) and South China Seas (SCS) by using 16S rRNA gene sequence analysis. The results demonstrated that the patterns of archaeal diversity in China Seas with the highest diversity in estuarine regions, moderate diversity in offshore regions and the lowest diversity in inshore regions. Phylogenetic analyses indicated that all archaeal 16S rRNA gene sequences obtained from China Seas belonged to the phyla *Crenarchaeota* and *Euryarchaeota*. In the estuarine areas, some archaeal clones were non-marine planktonic archaea, probably of terrestrial origin; while planktonic archaea dominated in inshore and offshore regions. Moreover, the dominant archaeal species were different from previous studies in other oceans of the world, revealing local specific characteristics of archaeal diversity.
2. The community structures of archaea were determined for an 11 m-long

subseafloor sediment core from the SCS. Our results revealed that the dominant phylogenetic groups of *Crenarchaeota* and *Euryarchaeota* are ubiquitous in pelagic sediments. RFLP types based clustering analysis and RDA analysis revealed depth-related changes in the community compositions of archaea in this sediment. Based on PCR screening of the functional genes of archaea, archaeal *amoA* genes were present in the surface layer of this sediment. Since our lab have recovered some 16S rRNA gene sequences of anammox- and ammonia-oxidizing bacteria from the same sample, we concluded that MGI *Crenarchaeota* together with other nitrifying microbes play important roles in the N cycling in the SCS sediments.

3. The community structure and ecological function of planktonic archaea were investigated for three pelagic stations in the SCS by using a combination of archaeal 16S rRNA, *amoA* and *accA* gene sequence analysis. Our results indicated that the abundance and genetic diversity of these genes were similar among the three stations. However, they are vertically stratified and significantly less abundant in the bathypelagia (< 1000 m) than in the epi- and mesopelagia at each station, except that archaeal *accA* genes were almost absent in the epipelagia. Furthermore, ratios of the archaeal *amoA* gene to 16S rRNA gene abundance slightly decreased from the epi- to bathypelagia, whereas the ratios of archaeal *accA* gene to MGI 16S rRNA gene or to archaeal *amoA* gene abundance increased with depth, suggesting that the trophic strategy of *Crenarchaeota* may change from the epi- to meso- or bathypelagia. Overall, our results show strong depth partitioning of crenarchaeal populations in the SCS and suggest a shift in their physiology and ecological function with increasing depth.
4. We studied the niche-partitioning and environmental adaptation mechanisms of marine *Crenarchaeota* in two pelagic stations of the ECS by using a combination of archaeal 16S rRNA, *amoA* and *accA* gene sequence analysis. Consistent with the results of our previous SCS investigation, we found most if not all *Crenarchaeota* contained *amoA* gene, but lack of *accA* gene, suggesting marine *Crenarchaeota* may live by chemolithoheterotrophy in the upper ocean. The results of ecotype stimulation showed that marine *Crenarchaeota* divided into several ecotypes to adapt local environments in the vertical scale of the upper ocean, implying that light may play an important role in controlling the niche-partitioning of marine *Crenarchaeota*.

5. Community structures of planktonic AOA and ammonia-oxidizing bacteria (AOB) were investigated for five high-altitude Tibetan lakes. Phylogenetic and statistical analyses indicated that a heterogeneous distribution of the AOA communities among lakes largely caused by salinity, whereas AOB communities were dominated by a unique monophylogenetic lineage within *Nitrosomonas*. Unlike depth-stratified community of marine *Crenarchaeota*, the uniform chemical properties within Tibetan lakes and their geographical isolation may favour relatively homogeneous AOA communities within lakes.

Keywords: planktonic archaea; *Crenarchaeota*; ammonia-oxidizing archaea; 16S rRNA; *amoA*; *accA*; autotrophic carbon fixation; phylogenetic analysis; ecological function; environmental adaptation mechanisms; China Seas; Tibetan lakes

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”.

Fulltexts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.